

OXYTROL™ for the treatment of overactive bladder with symptoms of

- Urge urinary incontinence
 - Urgency
 - Frequency
- Thin, flexible, translucent patch applied *twice weekly* for continuous delivery of oxybutynin 3.9 mg/day through the skin

OXYTROL is contraindicated in patients with urinary retention, gastric retention, or uncontrolled narrow-angle glaucoma and in patients who are at risk for these conditions. OXYTROL is also contraindicated in patients who have demonstrated hypersensitivity to oxybutynin or other components of the product. OXYTROL should be administered with caution in the following patients: those with hepatic or renal impairment; clinically significant bladder outflow obstruction; gastrointestinal obstructive disorders because of the risk of gastric retention; patients with gastroesophageal reflux.

Please see full prescribing information for OXYTROL.

When to apply OXYTROL

Before you begin treatment, choose any 2-day combination to use as your guide to change your patch (for example, every Sunday and Wednesday or Monday and Thursday). This will become your dosing schedule.

- Remind patients to change their patch on the *same 2 days* each week

Where to apply OXYTROL

Choose an area on the abdomen, hip, or buttock that is clean, dry, and smooth. The skin should have no cuts or irritations and should be fold-free.

- Avoid sun exposure; keep patch beneath clothing
- Avoid placing patch under waistband
- Skin should be free of body powder, lotion, and oil




abdomen



hip



buttock



In contact with skin, an OXYTROL patch delivers oxybutynin through the skin and into the bloodstream.

- Sunday/Wednesday
- Monday/Thursday
- Tuesday/Friday
- Wednesday/Saturday
- Thursday/Sunday
- Friday/Monday
- Saturday/Tuesday




How to apply OXYTROL™

Step 1.

Open the pouch by tearing along the arrows.

- Apply patch immediately. Do not store outside of sealed pouch



Step 2.

Apply one half of the patch to the skin.

- Remove one half of protective liner
- Press patch, adhesive side down, firmly onto skin



Step 3.

Apply the second half of the patch to the skin.

- Bend patch in half and gently roll remaining part onto skin
 - Liner should move off the patch
- Press the entire patch firmly onto skin



To remove the patch

- Remove slowly, fold in half (sticky sides together), and discard
- Gentle cleansing with warm water and mild soap should remove any adhesive residue
 - A small amount of baby oil may be used to remove excess residue

